OPEN ACCESS International Journal of Molecular Sciences ISSN 1422-0067 www.mdpi.com/journal/ijms

Review

A Review of Microsatellite Markers and Their Applications in Rice Breeding Programs to Improve Blast Disease Resistance

Gous Miah¹, Mohd Y. Rafii^{1,2,*}, Mohd R. Ismail^{1,2}, Adam B. Puteh², Harun A. Rahim³, Kh. Nurul Islam⁴ and Mohammad Abdul Latif^{2,5}

- ¹ Laboratory of Food Crops, Institute of Tropical Agriculture, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia; E-Mails: g_miah@yahoo.co.uk (G.M.); razi@upm.edu.my (M.R.I.)
- ² Department of Crop Science, Faculty of Agriculture, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia; E-Mails: adam@upm.edu.my (A.B.P.); alatif1965@yahoo.com (M.A.L.)
- ³ Agrotechnology and Bioscience Division, Malaysian Nuclear Agency, 43000 Kajang, Selangor, Malaysia; E-Mail: rahim@nuclearmalaysia.gov.my
- ⁴ Laboratory of Anatomy and Histology, Department of Veterinary Preclinical Sciences, Faculty of Veterinary Medicine, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia; E-Mail: kislam_2013@yahoo.com
- ⁵ Bangladesh Rice Research Institute, Gazipur 1701, Bangladesh
- * Author to whom correspondence should be addressed; E-Mail: mrafii@upm.edu.my; Tel.: +603-8947-1149.

Received: 12 August 2013; in revised form: 26 September 2013 / Accepted: 16 October 2013 / Published: 14 November 2013

Abstract: Over the last few decades, the use of molecular markers has played an increasing role in rice breeding and genetics. Of the different types of molecular markers, microsatellites have been utilized most extensively, because they can be readily amplified by PCR and the large amount of allelic variation at each locus. Microsatellites are also known as simple sequence repeats (SSR), and they are typically composed of 1–6 nucleotide repeats. These markers are abundant, distributed throughout the genome and are highly polymorphic compared with other genetic markers, as well as being species-specific and co-dominant. For these reasons, they have become increasingly important genetic markers in rice breeding programs. The evolution of new biotypes of pests and diseases as well as the pressures of climate change pose serious challenges to rice breeders, who would like to increase rice production by introducing resistance to multiple biotic and abiotic stresses.

Recent advances in rice genomics have now made it possible to identify and map a number of genes through linkage to existing DNA markers. Among the more noteworthy examples of genes that have been tightly linked to molecular markers in rice are those that confer resistance or tolerance to blast. Therefore, in combination with conventional breeding approaches, marker-assisted selection (MAS) can be used to monitor the presence or lack of these genes in breeding populations. For example, marker-assisted backcross breeding has been used to integrate important genes with significant biological effects into a number of commonly grown rice varieties. The use of cost-effective, finely mapped microsatellite markers and MAS strategies should provide opportunities for breeders to develop high-yield, blast resistance rice cultivars. The aim of this review is to summarize the current knowledge concerning the linkage of microsatellite markers to rice blast resistance genes, as well as to explore the use of MAS in rice breeding programs aimed at improving blast resistance in this species. We also discuss the various advantages, disadvantages and uses of microsatellite markers relative to other molecular marker types.

Keywords: simple sequence repeats; marker development and application; blast resistance; marker assisted selection; rice breeding

1. Introduction

Rice blast, which is caused by *Pyricularia grisea* (Cooke) Sacc., the anamorphous state of *Magnaporthe grisea* (T.T. Hebert) Barr [1], is the primary limiting biotic factor for rice production throughout the world. The use of resistant cultivars is the most effective and economical way to control rice blast disease, and therefore, breeding efforts to develop new resistant cultivars continue to be a priority for rice breeding programs. One of the challenges facing breeders during the development of improved rice cultivars, be they for conventional or organic agriculture, is the incorporation of disease resistance. Since the idea of indirect selection using genetic markers was first reported by Sax [2] over 80 years ago, and particularly in the last few decades, new technologies have emerged that allow breeders to more easily select changes at the DNA level. Much of the progress to date has centered on marker-assisted backcrossing or the pyramiding of genes against rice blast [3]. Presently, the integration of genomics and molecular-based breeding strategies for developing disease resistance, with gene-based marker assisted selection (MAS) being particularly effective, is a powerful method for efficient selection. In this context, pyramiding several major resistance genes into a valuable genetic background is simplified by the use of marker-based selection [4]. In short, the development and use of DNA markers has irrevocably changed the fields of rice genetics and breeding.

Molecular markers are now widely used to track loci and genome regions in crop breeding programs, as large numbers of molecular markers that are tightly linked to disease resistance traits are available in most major crop species [5–7]. The majority of molecular markers have been isolated from genomic DNA libraries or from libraries of randomly amplified PCR fragments. Molecular markers are essential for mapping genes of interest, marker-assisted breeding, and cloning genes using mapping-based cloning strategies [8]. Other uses of molecular markers include gene introgression

through backcrossing, germplasm characterization and phylogenetic analysis [6]. Of the various classes of existing markers, microsatellites have emerged as the markers of choice for plant breeding applications [9]. Restriction fragment length polymorphism (RFLP) analyses are not easily scalable to high-throughput methods, and random amplification of polymorphic DNA (RAPD) assays are often not reproducible or transferable between laboratories. Although both microsatellites and amplified fragment length polymorphisms (AFLPs) can both be used to efficiently identify polymorphisms, microsatellite-based methods are more readily automated [10]. In addition, AFLP analysis is not always straightforward, as seemingly individual bands may actually be composed of multiple fragments [11], particularly when using large genomic templates. Furthermore, molecular markers can be used to estimate overall genetic variability, determine the proportion of a genome that has been introgressed from a donor, identify genes that are phenotypically related to a particular analyzed trait, and select for traits during multiple rounds of introgression [12].

Identifying resistance genes using molecular markers is the basic prerequisite for performing MAS in resistance breeding programs [13]. PCR-based microsatellite markers have attracted a great deal of attention due to several key advantages. Most importantly, they are hypervariable, abundant and well distributed throughout the rice genome. Furthermore, these markers are readily accessible through published linkage maps and public databases, and they permit the differentiation between homozygous and heterozygous individuals [14,15]. Overall, the speed, reliability and cost-effectiveness of acrylamide gel-based microsatellite analysis make this method an attractive tool for MAS in blast resistance breeding programs [16].

In recent years, microsatellite markers have been widely used to screen, characterize and evaluate genetic diversity in cereal species [17]. In particular, microsatellite-based methods offer an attractive high-throughput and non-labor-intensive way to tag blast resistance genes in breeding programs. A number of microsatellite markers have been developed from publicly available databases (http://www.gramene.org) that are tightly linked with the Pi-ta², Pi-k^h and Pi-b resistance genes [18,19]. Furthermore, the development of molecular methods to efficiently identify novel resistance genes has the potential to greatly improve modern cultivars, and such methods would help accelerate the application of MAS and marker-assisted backcross (MAB) breeding in rice improvement programs. Use of these markers should also facilitate the development of multiline cultivars that carry one or more blast resistance genes, without the need for pathogenic inoculation or phenotyping. Microsatellite markers have been integrated into the molecular genetic maps of a number of plant species, and they have been successfully used to perform gene-mapping, population and evolutionary studies for the purpose of cultivar development. In this review, we summarize the known microsatellite markers that are linked to rice blast disease resistance. Furthermore, we investigate the organization of microsatellites within the rice genome and evaluate their usefulness as genetic markers. In particular, this review focuses on the availability of DNA markers linked to blast resistance in rice improvement programs and their potential use in MAS. The main objectives of this review are the following: (i) to help increase the efficiency of MAS in breeding-program crosses, thus improving the chances of developing new lines for commercial release; (ii) to review the basic principles and characteristics of commonly used microsatellite markers; (iii) to outline the advantages and limitations of these markers; and (iv) to provide examples of how microsatellites have been used in molecular breeding programs for blast resistance.

2. Microsatellites

The term microsatellite was first coined by Litt and Luty [20]. Microsatellites are simple repeated motifs consisting of 1 to 6 base pairs, and they can be found in both coding and non-coding regions. The mutation rate of this type of genetic marker has been estimated to be between 10^{-2} and 10^{-4} per generation. The primary advantage of microsatellites as genetic markers is that they are inherited in a Mendelian fashion as codominant markers. Furthermore, high polymorphism rates, high abundance and a broad distribution throughout the genome have made microsatellites one of the most popular genetic markers for use in plant breeding programs [21,22]. However, significant drawbacks do exist with respect to using microsatellite-based methods, including relatively high development costs and technical challenges during the construction of enriched libraries and species-specific primers.

3. Microsatellites: The Marker of Choice

The underlying principles and methodologies of various types of molecular markers that have been used in plants are shown in Table 1. The main challenge for researchers lies in selecting one or more of these markers for their specific purposes. The ideal type of genetic marker should be highly polymorphic, show codominant inheritance and be evenly distributed throughout the genome. In addition, particular marker sequences should be easy to access, and analyses should be low cost, high-throughput, reproducible, and transferable between laboratories, populations and/or species. Unfortunately, no marker type currently exists that meet all these requirements. However, based on the particular type of study, one can still choose among the different molecular marker systems to find the one that best suits your needs. A number of factors should be considered when choosing between the various molecular markers:

- (a) Marker system availability
- (b) Complexity of the technique and time investment
- (c) Estimated polymorphism levels within the study population
- (d) Quantity and quality of available DNA available
- (e) Transferability between laboratories, populations, pedigrees and species
- (f) The size and structure of the population to be studied
- (g) Availability of skilled workers and equipment
- (h) Cost per data-point and funding availability

(i) Method of marker inheritance (e.g., dominant *vs.* codominant) and the type of genetic information needed in the population [23–27].

In most areas of molecular genetics, microsatellites are now the marker of choice [28]. Microsatellites are also outstanding markers for fluorescent techniques, high-throughput analyses and multiplexing. However, high development costs and the significant effort required to design primer sets for a given study species remain limitations for using microsatellite markers to screen genomic libraries.

S.N.	Feature	RFLP	RAPD	AFLP	SSRs	SNPs
1	DNA Require (µg)	10	0.02	0.5-1.0	0.05	0.05
2	PCR based	No	Yes	Yes	Yes	Yes
3	DNA quality	High	High	Moderate	Moderate	High
4	No. of Polymorph loci analyzed	1–3	1.5-50	20-100	1–3	1
5	Type of polymorphism	Single base change,	Single base change,	Single base change,	Change in	Single nucleotide change,
		insertion, deletion	insertion, deletion	insertion, deletion	repeat length	insertion, deletion
6	Dominance	Co-dominant	Dominant	Dominant/Co-dominant	Co-dominant	Co-dominant
7	Reproducibility	High	Unreliable	High	High	High
8	Ease of use and development	Not easy	Easy	Easy	Easy	Easy
9	Automation	Low	Moderate	Moderate	High	High
10	Cost per analysis	High	Low	Moderate	Low	Low
11	Developmental cost	Low	Low	Moderate	High	High
12	Need for sequence data	Yes	No	No	Yes	Yes
13	Accuracy	Very high	Very low	Medium	High	Very high
14	Radioactive detection	Usually yes	No	No	No	Yes
15	Genomic abundance	High	Very high	Very high	Medium	Medium
16	Part of genome surveyed	Low copy coding regions	Whole genome	Whole genome	Whole genome	Whole genome
17	Level of polymorphism ^a	Low	Low to moderate	Low to moderate	High	High
18	Effective multiplex ratio ^b	Low	Medium	High	Medium	Medium
19	Marker index ^c	Low	Medium	High	Medium	Medium
20	Inheritance	Codominant	Dominant	Dominant	Codominant	Codominant
21	Detection of alleles	Yes	No	No	Yes	Yes
22	Utility for genetic mapping	Species specific	Cross specific	Cross specific	Species specific	Species specific
23	Utility in Marker assisted selection	Moderate	Low to moderate	Low to moderate	High	Low to moderate
24	Cost and labour involved in generation	High	Low-moderate	Low-moderate	High	High

Table 1. Important feature of different types of molecular markers.

^a Level of polymorphism (average heterozygosity) is an average of the probability that two alleles taken at random can be distinguished; ^b Effective multiplex ratio is the number of polymorphic loci analysed per experiment in the germplasm tested; ^c Marker index is the product of the average expected heterozygosity and the effective multiplex ratio. Source: [29–33].

4. Classification of Microsatellites

Microsatellites can be classified based on size, the nature of the repeated unit or their position within the genome. With respect to the number of nucleotides per repeat unit, microsatellites can be classified as mono-, di-, tri-, tetra-, penta- or hexa-nucleotide repeats (Table 2). Depending upon the arrangement of nucleotides within the repeat motifs, Weber [34] used the terms perfect, imperfect and compound to classify microsatellites, whereas Wang *et al.* [35] coined the terms simple perfect, simple imperfect, compound perfect and compound imperfect. Perfect repeats are tandem arrays of a single repeat motif, whereas imperfect repeats consist of otherwise perfect repeats that are sometimes interrupted by non-repeat sequences. In compound microsatellites, two basic repeat motifs are present together in various configurations. Jarne and Lagoda [36] coined the terms pure and interrupted to describe perfect or imperfect repeats, respectively, as shown in Table 2.

I ADIC 2. Classification of microsolemics.

(A) Based on the arrangement of nucleotides in the repeat motifs [34–36]				
Pure or perfect or simple perfect $(CA)n$ Simple imperfect $(AAC)n$ ACT $(AAC)n + 1$				
Compound or simple compound (CA)n (GA)n				
Interrupted or imperfect or compound imperfect (CCA) n TT (CGA) $n + 1$				
(B) Based on the number of nucleotides per repeat [31]				
Mononucleotide (A)n				
Dinucleotide (CA) <i>n</i>				
Trinucleotide (CGT)n				
Tetranucleotide (CAGA)n				
Pentanucleotide (AAATT)n				
Hexanucleotide (CTTTAA) n (n = number of variables)				
(C) Based on location of SSRs in the genome [31]				
Nuclear (nuSSRs)				
Chloroplastic (cpSSRs)				
Mitochondrial (mtSSRs)				

Table 3. Some potential benefits and weakness of the most commonly used markers.

Markers type	Benefits	Weakness
	-Co-dominant	-Need high-quality DNA
	-Genomic abundance high	-Laborious (compared to RAPD)
	-Highly reproducible	-Complex to automate
RFLP	-Better genome exposure	-Radioactive labeling essential
	-Applicable across the species	-Characterization of probe is essential
	-No need for sequence information	
	-Reliably used in plants	
	-Genomic abundance high	-No need of probe information
	-Better genome coverage	-Dominant markers
	-Sequence information unneeded	-Not reproducible
RAPD	-Perfect for automation	-Not suitable for across species
	-Requires less DNA	-Not well tested
	-No radioactive labeling	
	-More rapid	

22505

Markers type	Benefits	Weakness			
	-Easy to automate	-Not well-examined			
	-Genomic abundance high	-Cannot suitable across species			
	-Highly reproducible	-Sequence information needed			
SSR	-High polymorphism				
	-Multiple alleles				
	-Moderately genome coverage				
	-No radioactive labeling				
	-High polymorphism	-Very tricky due to changes in materials used			
	-Genomic abundance high	-Not reproducible			
AELD	-Can be used across species	-Very good primers needed			
ALL	-No need for sequence information				
	-Useful in preparing counting maps				
	-Works with smaller RFLP fragments				
	-Helpful in preparing counting maps	-Need sequence information			
Common to cool	-Highly reproducible	-Out of the target sites, mutation detection not possible			
sequence-tagged	-No radioactive labeling	-Laborious			
site (515)	-Can use filters many times	-Cloning and probe characterization required			
	-Moderate genome coverage				
	-Highly polymorphic	-Many informative bands per reaction			
Minisotallitas	-Multiallelic markers	-Band profiles can not be interpreted in terms of loci and alleles			
Minisatellites	-High reproducibility				
	-Low cost				
	Source: [20]				

Table 3. Cont.

Source: [29].

5. Comparative Advantages and Disadvantages of Microsatellite Markers

The use of molecular markers is still prohibitively expensive for most large-scale applications in rice breeding programs. Therefore, MAS methods are currently used for more targeted applications [37]. The advantages and disadvantages of the most commonly used markers are presented in Table 3.

6. Utility of Microsatellite Markers

A significant advance in the practical utilization of molecular markers was the development of microsatellite markers [38]. Microsatellites are extremely informative markers that can be used for a variety of population genetics studies. Microsatellites are also considered ideal markers for genetic mapping studies [36,39,40] and germplasm evolution. For instance, Sakai et al. [41] used microsatellite markers to investigate the genomic evolution of the African cultivated rice Oryza glaberrima and the Asian cultivated rice Oryza sativa, and they identified 2451 microsatellite from these two genomes. Of these 2451 microsatellites, 883 were identical among the three studied genomes (Oryza glaberrima, Oryza sativa indica and Oryza sativa japonica), whereas the remaining 1568 markers were variable. Microsatellite markers show a higher degree of polymorphism in rice compared with restriction fragment length polymorphisms (RFLPs), and they are also suitable for evaluating genetic diversity among closely related rice cultivars [42]. For MAS applications, microsatellite markers with high polymorphism information content (PIC) values are generally more useful. Parida et al. [43] observed

higher PIC in *indica* strains compared with aromatic, *japonica* and aus/wild strains, which agreed well with earlier observations using microsatellite and SNP markers [44–51]. Using their model, they were also able to identify four major genetically distinct groups within rice-indica, japonica, aromatics and aus/wild—based on population-structure analyses using microsatellites and SNP markers [45,48-50]. The estimated genetic diversity among the rice subpopulations in their study was significantly higher than previously observed using microsatellite and SNP markers [47,49] but was comparable to the diversity detected within a larger group of rice genotypes analyzed using microsatellite markers [51]. There exist several other examples of using microsatellites for these types of studies. Another interesting application of microsatellites in rice breeding was described by Liu and Wu [52]. In a study by Edwards et al. [53], genotyping microarrays utilizing single feature polymorphisms (SFPs) were used to assess the genetic variability across 20 diverse O. sativa varieties representing five different subpopulations, as determined using STRUCTURE [54] analysis with 169 microsatellite markers [49]. The highest level of polymorphism (66.2%) was found between the temperate japonica and indica subpopulations. The lowest levels of polymorphism were found within the temperate japonica sub-population (10.4%), which is also the least diverse subpopulation according to microsatellite markers [49].

As the order of markers along chromosomes is conserved within species and generally conserved between related species, this information can be used to create linkage maps, which are becoming increasingly available and detailed for a large number of species. This information is available for many marker types, and microsatellite-based maps are particularly well suited for genotyping [55]. For example, by analyzing such mapping information, it is possible to determine the fates of distinct parts of the genome over the course of developing multi-line varieties and composite-crosses [56]. Summaries of the various applications of microsatellite-based markers in plants are presented in Table 4 and Figure 1. However, it should be noted that each type of microsatellite-based marker possesses its own set of advantages and disadvantages based on mode of inheritance, informativeness, reproducibility, or procedural complexity, as well as economic concerns, such as cost and labor time. Therefore, the decision concerning which type of marker to use should be carefully considered based on the nature of the particular research project.

F 4	Marker type				
Features	Microsatellite	ISSR	SAMPL		
Abundance	High	High	Medium/high		
Locus specifity	Yes	No	No		
Nature of polymorphism	Variation in repeat length/number of motifs	Base changes (insertions, deletions),	(insertions, deletions) variation in		
		variation in microsatellite repeat	SSR repeat length/number of motifs		
Level of Polymorphism	High/very high	High/medium	High		
Inheritance mode	Codominance	Dominance/codominance	Codominance/dominance		
Reproducibility	High	High/medium	High		
Sequence information required	Yes	No	No		
Technical demands	medium/low (except for library construction	low/medium	medium		
	and screening)				
Costs	Medium	Low	Medium		
Labor	High (a labor-consuming step of library	Low	Medium		
	construction and screening)				
Time	Usually a time-consuming step of library	Low	Medium		
	construction and screening is needed				
Main applications	Linkage mapping, studies on genetic diversity,	Identification of cultivars,	Studies on genetic diversity,		
	gene tagging	phylogenetic studies	linkage Mapping		
Main advantages	High level of polymorphisms (up to 26 alleles),	Multilocus and highly polymorphic	Amplification of many		
	co-dominant mode of inheritance, very high	pattern production per reaction,	informative bands per reaction,		
	reproducibility	technical simplicity, low expenses	high reproducibility		
Problems	Frequently a small number of potential	Band profiles cannot be interpreted in	Relatively time consuming and		
	microsatellite loci are identified, polymerase	terms of loci and alleles, dominance of	labor-intensive procedure, high		
	slippage when analysing mono- and	alleles (frequently), similar-sized	complexity of amplification profiles		
	di-ucleotide repeats, co-migrating fragments	fragments may not be homologous	may occur		
	not always are homologous				

Table 4. A comparison of the main features of microsatellite-based marke
--

Source: [57].



Figure 1. Development and applications of microsatellite markers at a glance [31].

7. Microsatellite Marker Development

Completely sequenced genomes provide the basis upon which to design a large number of gene-based microsatellite markers. For example, rice (Oryza sativa L.) was the first cereal to have its genome completely sequenced, which has enabled the development of a large number of microsatellite markers [58]. Recently, Zhang et al. [59] developed 52,485 microsatellite markers that are polymorphic between *indica* and *japonica*. However, the difficulty now lies in choosing the most useful and informative microsatellite markers from such large datasets to use in rice genotyping applications. This problem can be overcome by constructing smaller, informative microsatellite marker databases composed of markers located in potentially functional genic sequences with relatively high polymorphic potential. Considering the excellent genetic attributes and higher predicted informativeness of genic non-coding microsatellite (GNMS) markers, Parida et al. [60] identified 19,555 perfect GNMS repeats on chromosomes 1 and 12 in rice. With the entire rice genome now sequenced, microsatellite markers can be developed within a few thousand base pairs of any gene. For example, a study by Goff et al. [61] suggests the presence, on average, of one microsatellite repeat (defined as at least eight repeats of a 2-4 bp motif) every 8 kb, yielding a total of 48,351 markers in the entire genome. Of course, not all these repeats can be developed into microsatellite markers [37], which can also be identified by screening database sequences or by screening libraries of clones (Figure 1).

PCR primers for amplifying microsatellite markers were designed by Rozen and Skaletsky [62] using the online program Primer3 by subjectively choosing primers flanking the repeat regions. The primers were then ordered from various commercial vendors and tested for their ability to amplify the microsatellites and differentiate polymorphisms among the parental lines used in the *Pi-z* mapping studies. Markers RM527 and RM6836, which were previously localized near the *Pi-z* locus [63], were obtained from the gramene website (http://www.gramene.org/) [64] and also included in primer testing [65].

Next-generation sequencing technologies (HighSSR, Roche 454 GS FLX) are now being used for microsatellite discovery with significant savings in cost and time [66–70]. Rapid progress in DNA sequencing technologies has substantially reduced costs while exponentially increasing throughput and accuracy. Currently, the most cost effective next-generation sequencing platform is the IlluminaHiSeq2000 [71], which can reduce costs 3400-fold relative to traditional sequencing methods; it is reasonable to expect that continued improvements will lead to even lower costs [72].

Fjellstrom et al. [73] designed DNA markers using four different methods. Five of these markers (RM101, RM138, RM144, RM155, and RM166) were based on an earlier set of microsatellites identified at Texas A & M University identified by screening the NCBI public DNA sequence database for repeated sequences, as described in Temnykh et al. [42]. The genomic locations of Pi-b, *Pi-k*, and *Pi-ta*² and their actual linkages were confirmed by mapping these genes in several of the populations described by Conaway et al. [19]. By mapping these initial markers relative to microsatellite markers developed using traditional methods at Cornell University [42], three additional tightly linked markers were subsequently identified: RM208, RM224, and RM266. After identifying candidate microsatellite markers from public database resources, the authors mapped these markers near to the blast resistance genes *Pi-b*, *Pi-k*, and *Pi-ta*² on rice chromosomes 2, 11 and 12, respectively. The public release of the Monsanto rice microsatellite database then allowed for the development of two additional linked microsatellite markers, which released as RM1233 and RM7102 [14]. Although DNA markers for rice blast resistance have been developed, most are not suitable for routine use in the MAS program involving large numbers of progeny. A dominant marker for the *Pi-b* gene, *Pibdom*, has also been developed based on the sequence of the cloned *Pi-b* gene [74] (GenBank accession AB013448). These markers should facilitate the introgression and pyramiding of these three blast resistance genes into new rice cultivars and elite lines [73]. In addition, a high-density microsatellite map with a genome coverage of approximately one microsatellite per 0.5 cM has been developed by the International Rice Microsatellite Initiative (IRMI) [75], which can be used for developing tightly linked markers for a variety agronomic traits, including blast resistance. Accessibility to the complete genomic sequences of the rice subspecies *indica* and *japonica* under public domain (http://rgp.dna.affrc.go.jp; http://www.genomics.org.cn) has enabled rice researchers to generate additional markers for the fine-scale mapping of targeted genes. For the purposes of obtaining a high-density linkage map for fine-scale mapping within their target region, new SSR, InDel, and CAPS markers were developed [76] using the publicly available rice genome sequence (http://rgp.dna.affrc.go.jp).

In species with large genomes, the conversion of microsatellite-containing sequences into useful markers can be difficult [77–80]. The recovery percentage of functional microsatellite primers in such genomes is usually low, which can be due to (i) the amplification of complex, weak or nonspecific patterns; (ii) lack of amplification; or (iii) non-polymorphic PCR products. Therefore, researchers

often choose to use tri- and tetra-nucleotide repeat motifs, as opposed to di-nucleotide motifs, as these generally yield fewer "stutter bands" [40,81]. However, di-nucleotide motifs are more common than tri- or tetra-nucleotide motifs and are therefore easier to use in combinatorial screens. The wide variety of DNA markers discovered and developed in rice is shown in Table 5.

Crop	Genome size (MB)	AFLP	SSR	SNP		
Rice	415–460 ^b	3,553 ^b	133 ^b	1,062 ^b	12,992 ^b	5,418,373 ^a
^a unuu nahi nim nih gau ^b Gramana wah brawaar (http://unuu gramana arg). Sauraa: [82]						

Table 5. Abundance of DNA markers discovered and developed in rice.

^a www.ncbi.nlm.nih.gov; ^b Gramene web browser (http://www.gramene.org). Source: [82].

Although microsatellite marker are considered to be the most practical genetic markers, their application remains somewhat limited due to the time and effort needed to develop them. Two general strategies are used to identify and create microsatellite markers: (i) searching for sequences containing microsatellites within databases; and (ii) constructing and screening genomic libraries with probes complementary to microsatellite sequences. Two websites recommended by Romero *et al.* [82] are shown in Table 6.

Table 6. Recommended	websites	for microsa	atellite markers.
----------------------	----------	-------------	-------------------

Gramene web browser	Gramene is a data resource for comparative genome analysis in the
http://www.gramene.org	grasses, in particular the cereals: rice, maize, oats etc. It provides
	comprehensive and in-depth information regarding markers used
	for mapping plant species such as RAPD, SSR, AFLP and RFLP.
MSU rice genome annotation projct	This website provides genome sequence from the Nipponbare
http://rice.plantbiology.msu.edu	subspecies of rice and annotation of the 12 rice chromosomes.
	0 [00]

Source: [82].

8. Abundance of Microsatellite Motifs in Rice

The frequency of microsatellite motifs varies significantly among different organisms [39,83,84]. The most abundant microsatellite motif reported in plants is (AT)n, whereas (AC)n is most abundant motif in the human genome. Currently, there are no reliable estimates of the number of (AT)n or (GC)n sites in rice due to the difficulty of the hybridization-based screening methods used to detect these motifs [85,86]. The size of the rice genome is $\sim 0.45 \times 10^9$ bp [87], whereas the size of the human genome is ~ 6.6 times larger $(3.0 \times 10^9 \text{ bp})$ [88]. These figures suggest that there should be one (AC)n site approximately every 360–450 kb in rice, compared with one every 40–80 kb in humans; similarly, it is estimated that there is one (GA)n motif every 225–330 kb in rice [85,86]. Three hundred and twenty three microsatellite markers identified by library screening [42,86,89,90] and GenBank searches of rice sequences [42,83,86] have been localized on the rice genetic map. The vast majority of these markers contain di- and tri-nucleotide motifs, with only seven loci containing tetra-nucleotide repeats, such as (AATT)n, (TTCC)n, (GATG)n, (ATGT)n, (GTAT)n, (ATTT)n, and (TTTG)n [42,86,89,90]. (GATA)n sequences, which to date have not been mapped in rice, are the most common tetra-nucleotide repeat, although only 270 of these motifs are found in the entire rice genome [91]. However, it should be kept in mind that it is the degree of polymorphism shown by microsatellites,

rather than their abundance within the genome, that ultimately determines their usefulness in genomic analyses [92].

The motif (CGG)*n* has been reported to be very abundant in rice and is interspersed throughout the genome [89]. Several studies based on GenBank searches of rice sequences identified a variety of microsatellites motifs [83,86,90]. In a study by Panaud *et al.* [85], 34-bp oligonucleotides (representing two di-nucleotide, seven tri-nucleotide, and four tetra-nucleotide motifs) were synthesized and used as probes for library screening. Screening of a 15-kb insert genomic library suggested that the relative frequency of the various microsatellite motifs was inversely correlated with the size of the repeat motif [38]. This finding was consistent with previous reports based on GenBank searches in a range of plant species [39]. The microsatellite (GATA)*n* has also been frequently used for DNA fingerprinting. For example, Davierwala *et al.* [92] identified three polymorphic (GATA)*n*-harboring loci (OS1A6, OS1H10 and OS2E7) that contained 7–13 repeat motifs by probing a genomic library from the cultivated rice strain *Oryza sativa* var. Basmati-370 with a oligonucleotide (GATA)4 probe.

Microsatellites can also be screened for using published rice sequences in the DDBJ databank, as numerous sequencing datasets, including those for EST sites, are available in this database. Homology searches can be performed to identify all di- and tri-nucleotide motifs, as well as four types of tetra-nucleotide repeats, and several of these are shown in Figure 2. Out of 11,798 total sequences, 369 contained complete repeats, and most types of repeat sequences, with the exception of (GC/CG)n, (AGT/TCA)n and (GACC/CTGG)n, were found. Sequences containing (CGG/GCC)n were most frequently identified in the published rice sequences, followed by sequences containing (GAG/CTC)n (Figure 2). Poly (CGG) loci were also abundant and were found throughout the rice genome [89], and the results presented here are likely representative of this feature of the rice genome. However, only a limited number of published rice sequences contain (AT/TA) sequences, although this motif is generally abundant in other plant genomes [83,84].

Figure 2. Frequencies of microsatellites in the rice sequences registered in the database [90].



9. Microsatellites for Tracking Blast Resistance in Rice

Many *Pi* genes confer resistance to overlapping spectra of blast pathotypes, and it is often difficult to monitor for the presence of individual resistance genes and pyramid these in breeding lines using

traditional phenotypic screening. Therefore, DNA markers provide a straightforward and rapid means to select for multiple blast resistance genes without performing extensive progeny testing or disease screening. DNA markers linked to several of the *Pi* genes have been localized on rice chromosomes, as well as markers for *Pi-ta* [93–95] and *Pi-b* [96]. Unfortunately, the majority of DNA markers for blast resistance are RFLPs, which are relatively labor intensive to analyze for use in breeding programs. Markers that can be analyzed by PCR are more amenable for breeding purposes, such as the ones developed for *Pi-2* [94] and *Pi-ta* [93].

The rice blast resistance gene *Pi-z*, which is present in the rice genotypes Zenith and Fukunishiki, represents a potential source of blast resistance for the northwestern Himalayan region of India. Sharma et al. [97] and Rathour et al. [98] both tested the reliability of microsatellite markers linked to Pi-z for assessing the blast resistance phenotype in commercially important crosses. A new set of microsatellite markers linked to Pi-z was developed by exploiting publicly available marker and genomic resources in rice. Of the three previously reported markers for Pi-z, only MRG5836 was found to be suitable for MAS. Among 17 microsatellites selected from the putative Pi-z locus, two (RM8225 and RM8226) co-segregated with MRG5836, and they were located at distance of 1.2-4.5 cM from the gene. In addition, a new microsatellite marker, SSR236, was derived from the (CT)16 repeat within the PAC clone P0502B12, and it showed even closer linkage to Pi-z. A survey of the allelic diversity at the loci containing the Pi-z-linked microsatellite markers revealed that the Fukunishiki- and Zenith-type alleles were not present in a majority of the local *indica* rice genotypes. Therefore, as these markers are polymorphic between the *Pi*-z donors and the great majority of local indica rice strains that were tested, they can be used as selection tools in rice breeding programs aimed at improving blast resistance in local rice varieties [99]. Conaway-Bormans et al. [63] identified three microsatellite markers that mapped at a distance of 0.0-11.5 cM from *Pi-z* in several different crosses segregating for the gene. Genetic distances between markers often vary in different mapping populations due to differences in the genetic backgrounds of the parental genotypes [100]. The gene *Pi-z* has been reported to be allelic with, or at least closely linked to, three other blast resistance genes, Pi-2, Pi-zt and Pi-9, which map close to the centromere of chromosome 6 [8,101]. Furthermore, the SSR236 marker has now bridged the 7.5-cM gap between the microsatellite markers RM8226 (54.1 cM) and RM3330 (61.6 cM) in the current version of the IRMI SSR map [99]. In addition, RM208 has been linked to Pi-b resistance, YL155 and YL183 have been linked to Pi-ta resistance in indica, and AP5659-1 has been linked to Pi-z resistance [65,73,102].

Gene name	Chromosome	Linked microsatellite marker	Rice variety	References
Pi-2(t)		RM140	Recombinant inbred lines	[103]
Pi33	8	RM72, RM44	IR64 × Azucena and Azucena × Bala	[104]
Pi-l(t)	11	RM1233*I and RM224	Near-isogenic lines C101LAC and C101A5	[16]
$Pi-k^h$	11	RM1233*I and RM206	Near-isogenic lines C101LAC and C101A5	[73]
Pi-k ^s	11	RM224	Near-isogenic lines C101LAC and C101A5	[73]
Pi37	1	RM140, RM302, RM212, FPSM1, FPSM2, FPSM4	-	[105]
Pi-b	2	RM166, RM138, RM208, RM266, RM138	Tohoku IL9 and Sasanishiki	[74]
Piz-t	6	RM225, RM226	Isogenic line C101A51 and cultivar CO39	[106]
Pi9	6	RM136	Cultivar TP309	[107]
Pid2	8	RM263	Variety LTH and Digu	[108]
Pi36	8	RM544	Q15 and Tsuyuake	[109]
Pita	12	OSM89, RM155, RM7102	Yashiro-mochi and Tsuyuake	[110]
Pi27(t)	1	RM151, RN259	Q14 and Q61	[111]
Pitp(t)	1	RM246	CO39 and Tetep	[112]
Pi35(t)	1	RM1216, RM1003	Hokkai 188 and Danghang-Shali	[113]
Pi37	1	RM302, RM212, FPSM1, FPSM2, FPSM4	C101PKT, CO39 and AS20-1 crossed with cultivar St. No. 1	[114]
Pid1(t)	2	RM262, RM208	Lijiangxintuanheigu (LTH) and Jiangnanxiangnuo (JNXN) crossed with Digu	[115]
Pig(t)	2	RM166	Q61 and Q14	[111]
Piyl	2	RM3248, RM20	Lijiangxintuanheigu (LTH) and Yanxian No.1	[116]
Piy2	2	RM3248, RM20	Lijiangxintuanheigu (LTH) and Yanxian No.1	[116]
Pi39	4	RM5473, RM3843	Mineasahi and Chubu 111	[117]
<i>Pi40(t)</i>	6	RM527, RM3330	Co39 and IR50 cross with IR65482-4-136-2-2	[118]
Pi36	8	RM5647	Aichi Asahi and Lijiangxintuanheigu (LTH) crossed with Q61	[119]
Pi38	11	RM206, RM21	CO39 and Tadukan	[120]

Table 7. Microsatellite markers linked to rice blast disease resistance gene.

 Table 7. Cont.

Gene name	Chromosome	Linked microsatellite marker	Rice variety	References
Pik-h	11	RM224, RM144, RM1233, RM144, RM1233,	HP2216 and Tetep	[73,121]
		RM224, RM206, TRS33, TRS26, RM144		
Pik-s	11	RM1233, RM224, RM144, RM1233,		[73]
		RM224, RM144	-	
Pita-2	12	OSM89, RM155, OSM89, RM7102,	Koshihikari cross with Fukunishiki (Piz+), Toride 1 (Piz-t+), K59	[122]
		OSM89, RM712	(Pit+), Kanto 51 (Pik+), Tsuyuake (Pik-m+), K60 (Pik-p+), BL 1	
			(Pib+), Yashiromochi (Pita+), and Pi No.4 (Pita-2+)	
Pi20(t)	12	RM1337, RM7102, RM54	Asominori and IR24	[123]
Pil5	9	RM316	Q61 and GA25	[124]
Pi36	8	RM5647-CRG2	Aichi Asahi and Lijiangxintuanheigu (LTH) cross with Q61	[119]
Pi37	1	RM543-FPSM1	cvs. C101PKT, CO39 and AS20-1 crossed with cultivar	[114]
			St. No. 1	
Pi39	12	RM27933-RM27940	Tsuyuake crossed with Q15	[109]
-	4	RM 5757	White Ponni x Moroberekan	[125]
-	4	RM 451	White Ponni x Moroberekan	[125]
-	2	RM 492	White Ponni x Moroberekan	[125]
-	2	RM208	Gulfmont*2/Te-Qing F12, Maybelle*2/Te-Qing F2	[73]
Pi - k^{s}	11	RM224	Maybelle*2/Kaybonnet F2, Maybelle*2/Lemont F2,	[73]
			Maybelle*2/Bengal F2, Maybelle*2/M-201 F2	
-	2	Pibdom	Gulfmont*2/Te-Qing F12	[73]
-	12	RM155	Maybelle*2/Kaybonnet F2	[73]
-	12	RM7102	Kaybonnet/M-204 F2	[73]
Pi40	6		wild Oryza species (O. australiensis)	[118]

Source: [126].

The *Pi20(t)* gene was identified from 160 Chinese *Magnaporthe oryzae* isolates, and among these, isolate 98095 can specifically differentiate the Pi20(t) gene present in cv. IR24. Two flanking and three co-segregating microsatellite markers for Pi20(t), which is located near the centromeric region of chromosome 12, were identified using 526 highly susceptible F_2 plants derived from a cross between Asominori (highly susceptible) and IR24 (resistant). The microsatellite OSR32 was mapped at a distance of 0.2 cM from Pi20(t), and the microsatellite RM28050 was mapped to the other side of Pi20(t) at a distance of 0.4 cM. The other three microsatellite markers, RM1337, RM5364 and RM7102, were observed to co-segregate with Pi20(t). In particular, RM1337 and RM5364 were found to be reliable markers of the resistance mediated by Pi20(t) in a wide range of elite rice germplasms from China. Therefore, these markers are useful tags for use in marker-assisted rice breeding programs aimed at incorporating Pi20(t) into advanced rice breeding lines [123]. Sharma et al. [127] concluded that RM25 and RM310 are two microsatellite markers linked to blast resistance in the Laxmi cultivar. These two markers are located 4.5 cM apart on chromosome 8 in rice [42]. An advanced backcross population of BC_3F_3 lines derived from the rice varieties Vandana and Moroberekan was analyzed for blast resistance by Wu et al. [128], and they identified four microsatellite markers (RM21, RM168, RM215 and RM250) that were significantly associated with the resistance gene.

This locus was mapped by Liu *et al.* [119] to a 5.8-cM interval bounded by RM5647 and RM8018 on the short arm of chromosome 8. This novel resistance gene has been tentatively designated as Pi36(t). According to a report by Liu, Pi-1 is located 6.8 cM away from the RM144 microsatellite; their results indicate that the physical distance between these two loci was between 57 and 72 kb [129]. Chen *et al.* [114] determined that the genetic distance between Pi-1 and the MRG4766 microsatellite marker was 1.3 cM. More recently, there has been a report concerning the identification of rice blast resistance using RM144 [130], indicating that it may be possible to identify this resistance gene using microsatellite markers. The known blast resistance genes and their linked microsatellite markers are shown in Table 7.

The Korean cultivar Suweon 365 carries three major resistance genes, Pi18, Pi21(t) and Pi22(t), that are effective against the Korean isolates KI-313, KJ-101, and KJ-201, respectively, and these genes have been tracked using microsatellite markers in an F₂ population derived from a cross of the Suweon and Chucheongbyeo cultivars [131]. Microsatellite markers were used to map the gene $Pi-k^h$, which confers resistance to blast races in the Himalayan region of Northeast India [121]. Pinheiro et al. [13] developed the cultivar Cica-8 by crossing the susceptible cultivar Metica-1 to the resistant cultivar Cica-8, and they found that one microsatellite marker, RM7102, was closely linked to the resistant allele. Yang et al. [132], while working on identifying and mapping the Pi41 gene, utilized resistant cultivar 93-11 and susceptible cultivar Nipponbare, as well as an F₂ population derived from a cross of these two cultivars. They studied 180 microsatellite markers and identified seven markers that were linked to Pi41, one of which was RM7102. The markers RM144, RM224 and yca72 are linked to the resistance genes *Pi-k^s*, *Pi1* and *Pia*, respectively [133,134]. Koide *et al.* [135] identified four microsatellite markers, RM7419, RM1268, RM6648 and RM5811 that were linked to Pish. With respect to Pib, both a gene-specific marker, Pibdom, and a co-segregation marker, RM208, have been reported [73]. Gouda et al. [136] found that the two microsatellite markers RM5926 and AP5659-5 were tightly linked to the Pi-1 and Piz-5 genes, respectively, in PRR78. Pan et al. reported that Pii(t), Pi3(t), Pi5(t) and Pi15(t) are all located within the same interval on chromosome 9 in

rice [124]. Moumeni and Leung reported that three microsatellite markers, RM224, RM179 and RM 277, on chromosomes 11 and 12 were tightly linked to components of rice blast resistance [137]. In addition, Liu *et al.* [109] revealed that RM247 and RM463 were located on chromosome 12 and were linked to the *Pi39* resistance gene. Abedi *et al.* [138] also identified four microsatellite markers, RM224, RM277, RM463 and RM179, which are linked to resistance genes on rice chromosomes in Iranian rice genotypes. This suggests there exists at least one race-specific resistance gene among the genetic sources of genotypes that confers resistance functions to the blast races. The marker AP5930, which is linked to *Piz-5* [65], and RM206, which is linked to *Pi54* [121], were used for foreground selection in both backcrossed and "selfed" generations. Finally, Singh *et al.* [139] also used a microsatellite marker (RM6100) when incorporating blast resistance into PRR78, an elite Basmati rice restorer line, through marker-assisted backcross breeding.

A large number of databases are available for selecting molecular markers linked to the Pi genes (http://www.gramene.org). Most of the known blast resistance genes have been mapped to chromosomes 12, 11 and 6 [140–143] through the work of a large number of researchers. For example, Pi-I(t), Pi-2(t) and Pi-4(t) were mapped to chromosomes 11, 6 and 12, respectively [144]. The microsatellite markers RM168, RM8225, RM1233, RM6836, RM5961 and RM413 were analyzed by Ashkani *et al.* [145], and they were linked to blast resistance genes specific to pathotype P7.2. Furthermore, molecular marker-assisted rice breeding programs have been developed with the aim of developing durable blast resistance in rice cultivars by pyramiding the resistance genes Pi-1(t), Pi-2(t) and Pi-33(t), which shows potential for controlling blast pathogen populations in Latin America [146].

10. Microsatellite Markers and Marker-Assisted Selection

The continued development of molecular markers promises to overcome most of the existing limitations associated with morphological markers. In particular, the tight linkage of a molecular marker to a gene can be exploited for indirect selection of traits in a breeding program, which is also referred to as MAS. Advances in the development of molecular markers and their implementation in cereal breeding programs have led to a greater understanding of rice genetics and genomes. Disease assays to evaluate resistance to rice blast are time-consuming and laborious procedures that require specialized facilities. However, PCR-based analyses promise to significantly reduce the amount of labor needed for evaluating phenotypes by prescreening with MAS. To accelerate the effectiveness of MAS, the map locations of target QTL must first be precisely determined, and several flanking markers must be developed [147]. Indeed, four QTLs derived from upland rice cultivars that control partial resistance to rice blast have been successfully pyramidized into lowland rice cultivars using MAS [147]. However, the successful application of MAS requires extremely tight linkages between markers and phenotypic traits. Some successful examples of using microsatellite markers in MAS to introduce blast resistance in rice are shown in Table 8.

,					
Table 8	Examples of MAS	application for	blast resistance	in rice.	

Application	Traits	Gene/QTLs	Markers used	References			
Gene surveys in	Blast disease	Pi-z	Microsatellite	[65]			
parental material							
Gene surveys in	Blast disease	Pi-ta	Gene-specific	[148]			
parental material			marker				
MAS applied for	Blast	Pil	Microsatellite	[149]			
backcross breeding			and ISSR				
Marker assisted	Submergence	Subchr9 QTL, Xa21, Bph and	Microsatellite	[150]			
backcrossing	tolerance, blast disease	blast QTLs, and quality loci	and STS				
	resistance, quality						
Marker assisted	Blast disease	-	Microsatellite	[151]			
backcrossing							
MAS applied for	Blast resistance BB	<i>Pi1</i> and <i>Pi2</i> for	Microsatellite	[152]			
backcross breeding		blast resistance					
Source [152 154]							

Source: [153,154].

Compared with previously reported RFLP markers linked to the Pi-1(t) gene [94,155], microsatellite markers are potentially more useful in developing countries where financial support is the principal limiting factor to establishing MAS rice breeding programs. Fuentes *et al.* [16] discussed the utility of DNA markers in MAS and gene pyramiding in rice breeding programs aimed at improving blast resistance. Indeed, PCR-based allele-specific markers provide an efficient system for MAS in blast resistance breeding programs [122]. However, the importance of using DNA markers within or flanking genes of interest during MAS strategies in rice should be stressed [73]. Finally, it is possible that several blast resistance genes could be combined using MAS in a single genetic background to develop rice cultivars with broad-spectrum durable resistance to blast [156].

11. Future Directions of Microsatellite Marker Research

Microsatellite markers provide an invaluable tool for plant geneticists and breeders, as detecting polymorphisms are a limiting factor in many breeding strategies. In the long term, the development of allele-specific markers for genes controlling disease resistance traits (e.g., blast disease resistance) will become increasingly important in the science of rice breeding. The choice of the most appropriate marker systems for a given program must be made on a case by case basis and will depend on many issues, including the availability of technology platforms, costs for marker development, species transferability, information content and ease of documentation. In addition, a higher degree of genetic variability and the localization of more markers on the rice linkage map will provide additional resources for genomic analysis and rice breeding. Therefore, there exist great opportunities for more efficient breeding programs and faster development times for new rice varieties resistant to biotic diseases in the future.

12. Conclusions

Molecular mapping of rice populations is a prerequisite to identifying markers closely linked to the desirable *Pi* resistance gene. In particular, microsatellite markers have become very important in rice breeding. Although many marker types exist within the rice genome, microsatellite markers are used in a wide range of studies due to their small size and repetitive nature, and they have played an important role in the identification of numerous important genetic loci. Microsatellite markers are also widely used in MAS programs to develop durably resistant cultivars against specific diseases. In recent years, the popularity of microsatellite-based markers has increased considerably. Microsatellites have been found to be highly polymorphic, genome-specific, abundant and co-dominant, and they have become important genetic markers in rice breeding programs for improving blast resistance.

This review has been specifically written for readers who want to use microsatellite markers for blast resistance improvement programs in rice cultivars, and will likely be useful for studies aimed at identifying the linkages between blast resistance genes and microsatellite markers. The techniques described will likely also prove useful for marker-assisted selection in the absence of appropriate pathogen isolates or when funds are limited, as is the case in most developing countries. Investigating the abundance and structure of rice microsatellite repeats, as well as their allelic variations and distributions should extend our knowledge concerning this class of tandem repeat in the rice genome. We hope that some of the ideas proposed in this article will encourage the rice scientific community to work together to convert rice from a model crop species into a model species for marker-assisted breeding.

Acknowledgments

The authors sincerely acknowledge the Long-term Research Grant Scheme (LRGS), Food Security Project, Ministry of Higher Education, Malaysia, for their financial support to conduct research activities on rice breeding as well as the ENSF for providing financial assistance.

Conflicts of Interest

The authors declare no conflict of interest.

References

- 1. Rossman, A.Y.; Howard, R.J.; Valent, B. *Pyricularia grisea*, the correct name for the rice blast fungus. *Mycologia* **1990**, *82*, 509–512.
- 2. Sax, K. The association of size differences with seed-coat pattern and pigmentation in *Phaseolus vulgaris. Genetics* **1923**, *8*, 552–560.
- Torres, A.M. Application of Molecular Markers for Breeding Disease Resistant Varieties in Crop Plants. In *Molecular Techniques in Crop Improvement*; Jain, S.M., Brar, D.S., Eds.; Springer Science and Business Media B.V.: Dordrecht, The Netherlands, 2010; pp. 185–205.
- 4. Song, W.Y.; Wang, G.L.; Zhu, L.H.; Fauquet, C.; Ronald, P. A receptor kinase-like protein encoded by the rice disease resistance gene, *Xa21*. *Science* **1995**, *270*, 1804–1806.
- Phillips, R.L.; Vasil, I.K. DNA-Based Markers in Plants. In DNA-Based Markers in Plants; Kluwer Academic Publishers: Dordrecht, The Netherlands, 2001; p. 497.

- 6. Jain, S.M.; Brar, D.S.; Ahloowalia, B.S. *Molecular Techniques in Crop Improvement*, 2nd ed.; Kluwer Academic Publishers: Dordrecht, The Netherlands, 2002; p. 772.
- 7. Gupta, P.K.; Varshney, R.K. Cereal Genomics: An Overview. In *Cereal Genomics*; Gupta, P.K., Varshney, R.K., Eds.; Kluwer Academic Press: Dordrecht, The Netherlands, 2004; p. 639.
- 8. Hayashi, K.; Hashimoto, N.; Daigen, M.; Ashikawa, I. Development of PCR-based SNP markers for rice blast resistance genes at the *Piz* locus. *Theor. Appl. Genet.* **2004**, *108*, 1212–1220.
- 9. Gupta, P.K.; Varshney, R.K. The development and use of microsatellite markers for genetics and plant breeding with emphasis on bread wheat. *Euphytica* **2000**, *113*, 163–185.
- 10. Shariflou, M.R.; Hassani, M.E.; Sharp, P.J. A PCR-based DNA marker for detection of mutant and normal alleles of the *Wx-D1* gene of wheat. *Plant Breed.* **2001**, *120*, 121–124.
- 11. Shan, X.; Blake, T.K.; Talbert, L.E. Conversion of AFLP markers to sequence-specific PCR markers in barley and wheat. *Theor. Appl. Genet.* **1999**, *98*, 1072–1078.
- 12. Brondani, C.; Hideo, P.; Rangel, N.; Borba, T.C.O.; Brondani, R.P.V. Transferability of micro-satellite and ISSR markers in Oryza species. *Hereditas* **2003**, *138*, 187–192.
- 13. Pinheiro, T.M.; de Araujo, L.G.; da Silva-Lobo, V.L.; Prabhu, A.S.; de Filippi, M.C. Tagging microsatellite marker to a blast resistance gene in the irrigated rice cultivar Cica-8. *Crop Breed. Appl. Biotechnol.* **2012**, *12*, 164–170.
- McCouch, S.R.; Teytelman, L.; Xu, Y.; Lobos, K.B.; Clare, K.; Walton, M.; Fu, B.; Maghirang, R.; Li, Z.; Xing, Y.; *et al.* Development and mapping of 2240 new SSR markers for rice (*Oryza sativa* L.). *DNA Res.* 2002, *9*, 199–207.
- Caixeta, E.T.; Oliveira, A.C.B.; Brito, G.G.; Sakiyama, N.S. Tipos de Marcadores Moleculares. In *Marcadores Moleculares*; Borem, A., Caixeta, E.T., Eds.; Editora UFV: Vicosa, Brazil, 2009; pp. 11–93.
- Fuentes, J.L.; Correa-Victoria, F.J.; Escobar, F.; Prado, G.; Aricapa, G.; Duque, M.C.; Thome, J. Identification of microsatellite markers linked to the blast resistance gene *Pi-1(t)* in rice. *Euphytica* 2008, *160*, 295–304.
- 17. Tautz, D. Hypervariability of simple sequences as a general source for polymorphic DNA markers. *Nucleic Acids Res.* **1989**, *17*, 6463–6471.
- Fjellstrom, R.; McClung, A.; Shank, A.R.; Marchetti, M.A.; Bormans, C.; Park, W.D. Progress on Development of Microsatellite Markers Associated with Rice Blast Resistance Genes. In Proceedings of the 29th Rice Technical Working Group Meeting, Little Rock, AR, USA, 24–27 February 2002; pp. 43–44.
- Conaway, C.; Cartinhour, S.; Ayres, N.; McClung, A.M.; Lai, X.H.; Marchetti, M.A.; Park, W.D. PCR Based Markers Linked to Blast Resistance Genes in Rice. In Proceedings of the 27th Rice Technical Working Group Meeting, Reno-Sparks, NV, USA, 1–4 March 1998; p. 77.
- 20. Litt, M.; Luty, J.A. A hypervariable microsatellite revealed by *in vitro* amplication of a dinucleotide repeat within the cardiac muscle actin gene. *Am. J. Hum. Genet.* **1989**, *44*, 397–401.
- 21. Morgante, M.; Hanafey, H.; Powell, W. Microsatellites are preferentially associated with nonrepetitive DNA in plant genome. *Nat. Genet.* **2002**, *30*, 194–200.
- 22. Wright, J.M.; Bentzen, P. Microsatellites: Genetic markers for the future. *Rev. Fish Biol. Fish.* **1994**, *4*, 384–388.

- 23. Staub, J.E.; Serquen, F.C. Genetic markers, map construction, and their application in plant breeding. *Hort Sci.* **1996**, *31*, 729–740.
- Karp, A.; Kresovich, S.; Bhat, K.V.; Ayad, W.G.; Hodgkin, T. Molecular Tools in Plant Genetic Resources Conservation: A Guide to the Technologies; IPGRI Technical Bulletin No. 2; International Plant Genetic Resources Institute (IPGRI): Rome, Italy, 1997.
- Wolfe, A.D.; Liston, A. Contribution of PCR-Based Methods to Plant Systematics and Evolutionary Biology. In *Molecular Systematics of Plants II: DNA Sequencing*; Soltis, D.E., Soltis, P.S., Doyle, J.J., Eds.; Kluwer Academic Publishers: Dordrecht, The Netherlands, 1998; pp. 43–86.
- 26. Mackay, T.F.C. The genetic architecture of quantitative traits. Ann. Rev. Genet. 2001, 35, 303–339.
- Rungis, D.; Hamberger, B.; Berube, Y.; Wilkin, J.; Bohlmann, J.; Ritland, K. Efficient genetic mapping of single nucleotide polymorphisms based upon DNA mismatch digestion. *Mol. Breed.* 2005, *16*, 261–270.
- 28. Gupta, P.K.; Varshney, R.K.; Sharma, P.C.; Ramesh, B. Molecular markers and their applications in wheat breeding. *Plant Breed.* **1999**, *118*, 369–390.
- 29. Kumar, P.; Gupta, V.K.; Misra, A.K.; Modi, D.R.; Pandey, B.K. Potential of molecular markers in plant biotechnology. *Plant Omics J.* **2009**, *2*, 141–162.
- 30. Benali, S.; Bencheikh, M.; Henni, J.E.; Claire, N. Advances of molecular markers application in plant pathology research. *Eur. J. Sci. Res.* **2011**, *50*, 110–123.
- 31. Kalia, R.K.; Manoj, K.R.; Sanjay, K.; Rohtas, S.; Dhawan, A.K. Microsatellite markers: An overview of the recent progress in plants. *Euphytica* **2011**, *177*, 309–334.
- 32. Semagn, K.; Bjornstad, A.; Ndjiondjop, M.N. An overview of molecular marker methods for plants. *Afr. J. Biotechnol.* **2006**, *5*, 2540–2568.
- 33. Farooq, S.; Azam, F. Molecular markers in plant breeding-II. Some pre-requisites for use. *Pak. J. Biol. Sci.* 2002, *5*, 1141–1147.
- 34. Weber, J.L. Informativeness of human (dC-dA)*n*(dGdT)*n* polymorphisms. *Genomics* **1990**, *7*, 524–530.
- 35. Wang, M.L.; Barkley, N.A.; Jenkins, T.M. Microsatellite markers in plants and insects. Part I. Applications of biotechnology. *Genes Genomes Genomics* **2009**, *3*, 54–67.
- 36. Jarne, P.; Lagoda, P.J.L. Microsatellites, from molecules to populations and back. *Trends Ecol. Evol.* **1996**, *11*, 424–429.
- 37. Mackill, D.J. Applications of genomics to rice breeding. IRRN 2003, 28, 9–15.
- 38. McCouch, S.R.; Chen, X.; Panaud, O.; Temnykh, S.; Xu, Y. Microsatellite marker development, mapping and applications in rice genetics and breeding. *Plant Mol. Biol.* **1997**, *35*, 89–99.
- 39. Morgante, M.; Olivieri, A.M. PCR-amplified microsatellites as markers in plant genetics. *Plant J.* **1993**, *1*, 175–182.
- 40. Hearne, C.M.; Ghosh, S.; Todd, J.A. Microsatellites for linkage analysis of genetic traits. *Tren Genet.* **1992**, *8*, 288–294.
- 41. Sakai, H.; Ikawa, H.; Tanaka, T.; *et al.* Distinct evolutionary patterns of *Oryza glaberrima* deciphered by genome sequencing and comparative analysis. *Plant J.* **2011**, *66*, 796–805.

- Temnykh, S.; Park, W.D.; Ayres, N.; Cartinhour, S.; Hauck, N.; Lipovtiesich, L. Mapping and genome organization of microsatellite sequences in rice (*Oryza sativa* L.). *Theor. Appl. Genet.* 2000, *100*, 697–712.
- Parida, S.K.; Mukerji, M.; Singh, A.K.; Singh, N.K.; Mohapatra T. SNPs in stress-responsive rice genes: Validation, genotyping, functional relevance and population structure. *BMC Genomics* 2012, 13, 426.
- 44. Rakshit, S.; Rakshit, A.; Matsumura, H.; Takahashi, Y.; Hasegawa, Y.; Ito, A.; Ishii, T.; Miyashita, T.; Terauchi, R. Large-scale DNA polymorphism study of *Oryza sativa* and *O. rufipogon* reveals the origin and divergence of Asian rice. *Theor. Appl. Genet.* **2007**, *114*, 731–743.
- Caicedo, A.L.; Williamson, S.H.; Hernandez, R.D.; Boyko, A.; Fledel-Alon, A.; York, T.L.; Polato, N.R.; Olsen, K.M.; Nielsen, R.; McCouch, S.R.; *et al.* Genome-wide patterns of nucleotide polymorphism in domesticated rice. *PLoS Genet.* 2007, *3*, e163.
- 46. Mather, K.A.; Caicedo, A.L.; Polato, N.R.; Olsen, K.M.; McCouch, S.; Purugganan, M.D. The extent of linkage disequilibrium in rice (*Oryza sativa* L.). *Genetics* **2007**, *177*, 2223–2232.
- 47. Agrama, H.A.; Eizenga, G.E. Molecular diversity and genome-wide linkage disequilibrium patterns in a worldwide collection of *Oryza sativa* and its wild relatives. *Euphytica* **2008**, *160*, 339–355.
- 48. Ni, J.; Colowitb, P.M.; Mackill, D.J. Evaluation of genetic diversity in rice subspecies using microsatellite markers. *Crop Sci.* **2002**, *42*, 601–607.
- 49. Garris, A.J.; Tai, T.H.; Coburn, J.; Kresovich, S.; McCouch, S.R. Genetic structure and diversity in *Oryza sativa* L. *Genetics* **2005**, *169*, 1631–1638.
- 50. Agrama, H.A.; Eizenga, G.C.; Yan, W. Association mapping of yield and its components in rice cultivars. *Mol. Breed.* 2007, *19*, 341–356.
- 51. Jin, L.; Lu, Y.; Xiao, P.; Sun, M.; Corke, H.; Bao, J. Genetic diversity and population structure of a diverse set of rice germplasm for association mapping. *Theor. Appl. Genet.* **2010**, *121*, 475–487.
- 52. Liu, X.C.; Wu, J.L. SSR heterotic patterns of parents for making and predicting heterosis. *Mol. Breed.* **1998**, *4*, 263–268.
- 53. Edwards, J.D.; Janda, J.; Sweeney, M.T.; Gaikwad, A.B.; Liu, B.; Leung, H.; Galbraith, D.W. Development and evaluation of a high-throughput, low-cost genotyping platform based on oligonucleotide microarrays in rice. *Plant Methods* **2008**, *4*, 13.
- 54. Falush, D.; Stephens, M.; Pritchard, J.K. Inference of population structure using multilocus genotype data: Linked loci and correlated allele frequencies. *Genetics* **2003**, *164*, 1567–1587.
- Karakousis, A.; Barr, A.R.; Chalmers, K.J.; Eckermann, P.J.; Holton, T.A.; Henry, R.J.; Lim, P.; Langridge, P. Potential of SSR markers for plant breeding and variety identification in Australian barley germplasm. *Aust. J. Plant Physiol.* 2003, *54*, 1197–1210.
- 56. Enjalbert, J.; Goldringer, I.; Paillard, S.; Brabant, P. Molecular markers to study genetic drift and selection in wheat populations. *J. Exp. Bot.* **1999**, *50*, 283–290.
- 57. Rakoczy-Trojanowska, M.; Bolibok, H. Characteristics and a comparison of three classes of microsatellite-based markers and their application in plants. *Cell Mol. Biol. Lett.* **2004**, *9*, 221–238.

- 58. International Rice Genome Sequencing Project. The map based sequence of rice genome. *Nature* **2005**, *436*, 793–800.
- 59. Zhang, Z.; Deng, Y.; Tan, J.; Hu, S.; Yu, J.; Xue, Q. A genome-wide microsatellite polymorphism database for the indica and japonica rice. *DNA Res.* **2007**, *14*, 37–45.
- 60. Parida, S.K.; Dalal, V.; Singh, A.K.; Singh, N.K.; Mohapatra, T. Genic non-coding microsatellites in the rice genome: Characterization, marker design and use in assessing genetic and evolutionary relationships among domesticated groups. *BMC Genomics* **2009**, *10*, 140.
- Goff, S.A.; Ricke, D.; Lan, T.H.; Presting, G.; Wang, R.; Dunn, M.; Glazebrook, J.; Sessions, A.; Oeller, P.; Varma, H.; *et al.* A draft sequence of the rice genome (*Oryza sativa* L. ssp. *japonica*). *Science* 2002, *296*, 92–100.
- Rozen, S.; Skaletsky, H.J. Primer3 on the WWW for General Users and for Biologist Programmers. In *Bioinformatics Methods and Protocols: Methods in Molecular Biology*; Misener, S., Krawetz, S.A., Eds.; Humana Press Inc.: Totowa, NJ, USA, 2000; pp. 365–386.
- Conaway-Bormans, C.A.; Marchetti, M.A;, Johnson, C.W.; McClung, A.M.; Park, W.D. Molecular markers linked to the blast resistance gene *Pi-z* in rice for use in marker-assisted selection. *Theor. Appl. Genet.* 2003, *107*, 1014–1020.
- Ware, D.; Jaiswal, P.; Ni, J.; Pan, X.; Chang, K.; Clark, K.; Teytelman, L.; Schmidt, S.; Zhao, W.; Cartinhour, S.; *et al.* Gramene: A resource for comparative grass genomics. *Nucleic Acids Res.* 2002, *30*, 103–105.
- Fjellstrom, R.; McClung, A.M.; Shank, A.R. SSR markers closely linked to the *Pi-z* locus are useful for selection of blast resistance in a broad array of rice germplasm. *Mol. Breed.* 2006, *17*, 149–157.
- Abdelkrim, J.; Robertson, B.; Stanton, J.A.; Gemmell, N. Fast, cost-effective development of species-specific microsatellite markers by genomic sequencing. *BioTechniques* 2009, 46, 185–192.
- Dutta, S.; Kumawat, G.; Singh, B.P.; Gupta, D.K.; Singh, S.; Dogra, V.; Gaikwad, K.; Sharma, T.R.; Raje, R.S.; Bandhopadhya, T.K.; *et al.* Development of genic-SSR markers by deep transcriptome sequencing in pigeonpea [*Cajanus cajan* (L.) Millspaugh]. *BMC Plant Biol.* 2011, *11*, 17.
- 68. Saarinen, E.V.; Austin, J.D. When technology meets conservation: Increased microsatellite marker production using 454 genome sequencing on the endangered Okaloosa darter (*Etheostoma okaloosae*). *J. Hered.* **2010**, *101*, 784–788.
- Santana, Q.; Coetzee, M.; Steenkamp, E.; Mlonyeni, O.; Hammond, G.; Wingfield, M.; Wingfield, B. Microsatellite discovery by deep sequencing of enriched genomic libraries. *Biotechniques* 2009, 46, 217–223.
- Churbanov, A.; Ryan, R.; Hasan, N.; Bailey, D.; Chen, H.; Milligan, B; Houde, P. HighSSR: High-throughput SSR characterization and locus development from next-gen sequencing data. *Bioinformatic* 2012, 28, 2797–2803.
- Liu, L.; Li, Y.; Li, S.; Hu, N.; He, Y.; Pong, R.; Lin, D.; Lu, L.; Law, M. Comparison of next-generation sequencing systems. *J. Biomed. Biotechnol.* 2012, doi:10.1155/2012/251364.
- Moran, C. Looking Back to Move Forward—A Personal Perspective on Pig Molecular Genetics from RFLPs to Nextgen Sequencing. In Proceedings of the AGBU Pig Genetics Workshop, Armidale, NSW, Australia, 24–25 October 2012; pp. 7–12.

- 73. Fjellstrom, R.; Conaway-Bormans, C.A.; McClung, A.M.; Marchetti, M.A.; Shank, A.R.; Park, W.D. Development of DNA markers suitable for marker assisted selection of three *Pi* genes conferring resistance to multiple *Pyricularia grisea* pathotypes. *Crop Sci.* 2004, *44*, 1790–1798.
- Wang, Z.X.; Yano, M.; Yamanouchi, U.; Iwamoto, M.; Monna, L.; Hayasaka, H.; Katayose, Y.; Sasaki, T. The *Pib* gene for disease resistance belongs to the nucleotide binding and leucine-rich repeat class of plant disease resistance genes. *Plant J.* 1999, *19*, 55–66.
- 75. McCouch, S.R. Development and mapping of 2240 new SSR markers for rice (*Oryza sativa* L.). *DNA Res.* **2002**, *9*, 193–207.
- Zhang, Y.X.; Wang, Q.; Jiang, L.; Liu, L.L.; Wang, B.X.; Shen, Y.Y.; Cheng, X.N.; Wan, J. Fine mapping of qSTV11KAS, a major QTL for rice stripe disease resistance. *Theor. Appl. Genet.* 2011, *122*, 1591–1604.
- 77. Song, Q.J.; Fickus, E.W.; Cregan, P.B. Characteristics of trinucleotide markers in wheat. *Theor. Appl. Genet.* **2002**, *104*, 286–293.
- 78. Pfeiffer, A.; Olivieri, A.M.; Morgante, M. Identification and characterization of microsatellites in Norway spruce (*Picea abies* K). *Genome* **1997**, *40*, 411–419.
- 79. Roder, M.S.; Plaschke, J.; Knig, S.U.; Berner, A.; Sorrels, M.E.; *et al.* Abundance, variability and chromosomal location of microsatellite in wheat. *Mol. Gen. Genet.* **1995**, *246*, 327–333.
- 80. Kostia, S.; Varvio, S.L.; Vakkari, P.; Pulkkinen, P. Microsatellite sequences in *Pinus sylvestris*. *Genome* **1995**, *38*, 1244–1248.
- 81. Diwan, N.; Cregan, P.B. Automated sizing of fluorescent labeled simple sequence repeat markers to assay genetic variation in soybean. *Theor. Appl. Genet.* **1997**, *95*, 723–733.
- 82. Romero, G.; Adeva, C.; Battad, Z., II. Genetic fingerprinting: Advancing the frontiers of crop biology research. *Philipp. Sci. Lett.* **2009**, *2*, 8–13.
- 83. Wang, Z.; Weber, J.L.; Zhong, G.; Tanksley, S.D. Survey of plant short tandem DNA repeats. *Theor. Appl. Genet.* **1994**, *88*, 1–6.
- 84. Lagercrantz, U.; Ellegren, H.; Andersson, L. The abundance of various polymorphic microsatellite motifs differs between plants and vertebrates. *Nucl Acids Res.* **1993**, *21*, 1111–1115.
- 85. Panaud, O.; Chen, X.; McCouch, S.R. Frequency of microsatellite sequences in rice (*Oryza sativa* L.). *Genome* **1995**, *38*, 1170–1176.
- 86. Wu, K.S.; Tanksley, S.D. Abundance, polymorphism and genetic mapping of microsatellites in rice. *Mol. Gen. Genet.* **1993**, *241*, 225–235.
- 87. Arumunagathan, K.; Earle, E.D. Nuclear DNA content of some important plant species. *Plant Mol. Biol.* **1991**, *9*, 208–218.
- 88. Morton, N.E. Parameters of the human genome. Proc. Natl. Acad. Sci. USA 1991, 88, 7474–7476.
- 89. Zhao, X.; Kochert, G. Characterization and genetic mapping of a short, highly repeated, interspersed DNA sequence from rice (*Oryza sativa* L.). *Mol. Gen. Genet.* **1992**, *231*, 353–359.
- 90. Akagi, H.; Yokozeki, Y.; Inagaki, A.; Fujimura, T. Microsatellite DNA markers for rice chromosomes. *Theor. Appl. Genet.* **1996**, *93*, 1071–1077.
- Squirrel, J.; Hollingsworth, P.M.; Woodhead, M.L.; Russell, J.; Lowe, A.J.; Gibby, M.; Powell, W. How much effort is required to isolate nuclear microsatellites from plants? *Mol. Ecol.* 2003, *12*, 1339–1348.

- 92. Davierwala, A.P.; Ramakrishna, W.; Chowdari, V.; Ranjekar, P.K.; Gupta, V.S. Potential of (GATA)*n* microsatellites from rice for inter- and intraspecific variability studies. *BMC Evolut. Biol.* 2001, *1*, 7.
- 93. Jia, Y.; Wang, Z.; Singh, P. Development of dominant rice blast *Pi-ta* resistance gene markers. *Crop Sci.* **2002**, *42*, 2145–2149.
- Hittalmani, S.; Parco, A.; Mew, T.V.; Ziegler, R.S.; Huang, N. Fine mapping and DNA marker-assisted pyramiding of the three major genes for blast resistance in rice. *Theor. Appl. Genet.* 2000, 100, 1121–1128.
- 95. Nakamura, S.; Asakawa, S.; Ohmido, N.; Fukui, K.; Shimizu, N.; Kawasaki, S. Construction of an 800-kb contig in the nearcentromeric region of the rice blast resistance gene *Pi-ta2* using a highly representative rice BAC library. *Mol. Gen. Genet.* **1997**, *254*, 611–620.
- 96. Monna, L.; Miyao, A.; Zhong, H.S.; Yano, M.; Iwamoto, M.; Umehara, Y.; Kurata, N.; Haysaka, H.; Sasaki, T. Saturation mapping with subclones of YACs: DNA marker production targeting the rice blast disease resistance gene, *Pi-b. Theor. Appl. Genet.* **1997**, *94*, 170–176.
- Sharma, T.R.; Chauhan, R.S.; Singh, B.M.; Sagar, V.; Paul, R.; Rathour, R. RAPD and virulence analyses of *Magnaporthe grisea* rice populations from north-western Himalayan region of India. *J. Phytopathol.* 2002, *150*, 649–656.
- 98. Rathour, R.; Singh, B.M.; Plaha, P. Virulence structure of *Magnaporthe grisea* rice population from north-western Himalayas. *Phytoparastica* **2006**, *34*, 281–291.
- 99. Rathour, R.; Chopra, M.; Sharma, T.R. Development and validation of microsatellite markers linked to the rice blast resistance gene *Pi-z* of Fukunishiki and Zenith. *Euphytica* **2008**, *163*, 275–282.
- 100. Wang, Z.; Taramino, G.; Yang, D.; Liu, G.; Tingey, S.V.; Miao, G.H.; Wang, G.L. Rice ESTs with disease-resistance gene or defense-response gene like sequences mapped to regions containing major resistance genes or QTLs. *Mol. Genet. Genomics* 2001, 265, 302–310.
- 101. Liu, G.; Lu, G.; Zeng, L.; Wang G.L. Two broad-spectrum blast resistance genes, *Pi9 (t)* and *Pi2 (t)*, are physically linked on rice chromosome 6. *Mol. Genet. Genomics* **2002**, *267*, 472–480.
- 102. Jia, Y.; Wang, Z.; Fjellstrom, R.G.; Moldenhauer, K.A.; Azam, M.A.; Correll, J.; Lee, F.N.; Xia Y.; Rutger, J.N. Rice *Pi-ta* gene confers resistance to the major pathotypes of the rice blast fungus in the United States. *Phytopathology* **2004**, *94*, 296–301.
- 103. Jiang, J.; Wang, S. Identification of a 118-kb DNA fragment containing the locus of blast resistance gene *Pi-2(t)* in rice. *Mol. Genet. Genomics* **2002**, *268*, 249–252.
- 104. Berruyer, R.; Adreit, H.; Milazzo, J.; Gaillard, S.; Berger, A.; Dioh, W.; Lebrun, M.H.; Tharreau, D. Identification and fine mapping of *Pi33*, the rice resistance gene corresponding to the *Magnaporthe grisea* avirulence gene *ACE1*. *Theor. Appl. Genet.* **2003**, *107*, 1139–1147.
- 105. Lin, F.; Chen, S.; Que, Z.; Wang, L.; Liu, X.; Pan, Q. The blast resistance gene *Pi37* encodes a nucleotide binding site leucine-rich repeat protein and is a member of a resistance gene cluster on rice chromosome 1. *Genetics* **2007**, *177*, 1871–1880.
- 106. Zhou, B.; Qu, S.; Liu, G.; Dolan, M.; Sakai, H.; Lu, G.; Bellizzi, M.; Wang, G.L. The eight amino acid differences within three leucine-rich repeats between *Pi2* and *Piz-t* resistance proteins determine the resistance specificity to *Magnaporthe grisea*. *Mol. Plant Microbe Interact.* 2006, 19, 1216–1228.

- 107. Qu, S.; Liu, G.; Zhou, B.; Bellizzi, M.; Zeng, L.; Dai, L.; Han, B.; Wang, G.L. The broad-spectrum blast resistance gene *Pi9* encodes a nucleotide binding site-leucine-rich repeat protein and is a member of a multigene family in rice. *Genetics* **2006**, *172*, 1901–1914.
- 108. Chen, X.; Shang, J.; Chen, D.; Lei, C.; Zou, Y.; Zhai, W.; *et al.* A Blectin receptor kinase gene conferring rice blast resistance. *Plant J.* **2006**, *46*, 794–804.
- 109. Liu, X.; Yang, Q.; Lin, F.; Hua, L.; Wang, C.; Wang, L.; Pan, Q. Identification and fine mapping of *Pi39(t)*, a major gene conferring the broad-spectrum resistance to *Magnaporthe oryzae*. *Mol. Genet. Genomics* 2007, 278, 403–410.
- 110. Bryan, G.T.; Wu, K.; Farrall, L.; Jia, Y.; Hershey, H.P.; McAdams, S.A.; Tarchini, R.; Donaldson, G.; Faulk, K.; Valent, B. A single amino acid difference distinguishes resistant and susceptible alleles of the rice blast resistance gene *Pi-ta*. *Plant Cell* **2000**, *12*, 2033–2045.
- Zhu, M.; Wang, L.; Pan, Q. Identification and characterization of a new blast resistance gene located on rice chromosome 1 through linkage and differential analyses. *Phytopathology* 2004, 94, 515–519.
- 112. Barman, S.R.; Gowda, M.; Venu, R.C.; Chattoo, B.B. Identification of a major blast resistance gene in the rice cultivar "Tetep". *Plant Bleed.* **2004**, *123*, 300–302.
- 113. Nguyen, T.T.T.; Koizumi, S.; La, T.N.; Zenbayashi, K.S.; Ashizawa, T.; Yasuda, N.; Imazaki, I.; Miyasaka, A. *Pi35(t)*, a new gene conferring partial resistance to leaf blast in the rice cultivar Hokkai 188. *Theor. Appl. Genet.* 2006, *113*, 697–704.
- Chen, S.; Wang, L.; Que, Z.; Pan, R.; Pan, Q. Genetic and physical mapping of *Pi37(t)*, a new gene conferring resistance to rice blast in the famous cultivar St. No. 1. *Theor. Appl. Genet.* 2005, *111*, 1563–1570.
- 115. Chen, X.W.; Li, S.G.; Xu, J.C.; Zhai, W.X.; Ling, Z.Z.; Ma, B.T.; Wang, Y.P.; Wang, W.M.; Cao, G.; Ma, Y.Q.; *et al.* Identification of two blast resistance genes in a rice variety, Digu. *J. Phytopathol.* 2004, 152, 77–85.
- 116. Lei, C.L.; Huang, D.Y.; Li, W.; Wang, J.L.; Liu, Z.L.; Wang, X.T.; Shi, K.; Cheng, Z.J.; Zhang, X.; Ling, Z.Z.; *et al.* Molecular mapping of a blast resistance gene in an *indica* rice cultivar Yanxian No. 1. *Rice Genet. Newsl.* 2005, *22*, 76–77.
- 117. Terashima, T.; Fukuoka, S.; Saka, N.; Kudo, S. Mapping of a blast field resistance gene *Pi39(t)* of elite rice strain Chubu 111. *Plant Breed*. **2008**, *127*, 485–489.
- 118. Jeung, J.U.; Kim, B.R.; Cho, Y.C.; Han, S.S.; Moon, H.P.; Lee, Y.T.; Jena, K.K. A novel gene, *Pi40(t)*, linked to the DNA markers derived from NBS-LRR motifs confers broad spectrum of blast resistance in rice. *Theor. Appl. Genet.* 2007, *115*, 1163–1177.
- Liu, X.Q.; Wang, L.; Chen, S.; Lin F.; Pan, Q.H. Genetic and physical mapping of *Pi36(t)*, a novel rice blast resistance gene located on rice chromosome 8. *Mol. Gen. Genomics* 2005, 274, 394–401.
- 120. Gowda, M.; Roy-Barman, S.; Chattoo, B.B. Molecular mapping of a novel blast resistance gene *Pi38* in rice using SSLP and AFLP markers. *Plant Breed*. **2006**, *125*, 596–599.
- 121. Sharma, T.R.; Madhav, M.S.; Singh, B.K.; Shanker, P.; Jana, T.K.; Dalal, V.; Pandit, A.; Singh, A.; Gaikwad, K.; Upreti, H.C.; *et al.* High resolution mapping, cloning and molecular characterization of the *Pi-k^h* gene of rice, which confers resistance to *M. grisea. Mol. Gen. Genom.* 2005, 274, 569–578.

- 122. Hayashi, K.; Yoshida, H.; Ashikawa, I. Development of PCR-based allele-specific and InDel marker sets for nine rice blast resistance genes. *Theor. Appl. Genet.* **2006**, *113*, 251–260.
- 123. Li, W.; Lei, C.; Cheng, Z.; Jia, Y.; Huang, D.; Wang, J.; Wang, J.; Zhang, X.; Su, N.; Guo, X.; *et al.* Identification of SSR markers for a broad-spectrum blast resistance gene *Pi20(t)* for marker-assisted breeding. *Mol. Breed.* 2008, *22*, 141–149.
- 124. Pan, Q.H.; Hu, Z.D.; Takatoshi, T.; Wang, L. Fine mapping of the blast resistance gene *Pi15*, linked to *Pii* on rice chromosome 9. *Acta Bot. Sin.* **2003**, *45*, 871–877.
- 125. Selvaraj, I.C.; Pothiraj, N.; Thiyagarajan, K.; Bharathi, M.; Rabindran, R. Identification of microsatellite (SSR) and RAPD markers linked to rice blast disease resistance gene in rice (*Oryza sativa* L.). *Afr. J. Biotechnol.* **2011**, *10*, 3301–3321.
- 126. Mackill, D.J.; Junjian, N. Molecular Mapping and Marker Assisted Selection for Major-Gene Traits in Rice. In *Rice Genetics IV*; Khush, G.S., Brar, D.S., Hardy, B., Eds.; International Rice Research Institute, Science Publishers, Inc.: Enfield, NH, USA, 2001; pp. 140–141.
- 127. Sharma, R.C.; Shrestha S.M.; Pandey, M.P. Inheritance of blast resistance and associated microsatellite markers in rice cultivar Laxmi. *J. Phytopathol.* **2007**, *155*, 749–753.
- 128. Wu, J.L.; Sinha, P.K.; Variar, M.; Zheng, K.L.; Leach, J.E.; Courtois, B.; Leung, H. Association between molecular markers and blast resistance in an advanced backcross population of rice. *Theor. Appl. Genet.* **2004**, *108*, 1024–1032.
- 129. Liu, S.; Xue, Y. Locating rice blast resistance gene by DNA micro-satellite markers. *J. Triple Gorge Univ.* **2003**, *6*, 574–576.
- Lin, H.; Jin, X.; Zhang, Y.; Wang, X. Identification of rice blast resistance gene *Pi-1* in some japonica rice varieties with SSR Marker. *J. Heilongjiang August First Land Reclam. Univ.* 2006, *II*, 13–15.
- 131. Ahn, S.N.; Kim, Y.K.; Hong, H.C.; Han, S.S.; Kwon, S.J.; Choi, H.C.; Moon, H.P.; McCouch, S.R. Molecular mapping of a new gene for resistance to rice blast (*Pyricularia grisea* Sacc.). *Euphytica* 2000, 116, 17–22.
- 132. Yang, Q.; Lin, F.; Wang, L.; Pan, Q. Identification and mapping of *Pi41*, a major gene c onferring resistance to rice blast in the *Oryza sativa* subsp. Indica reference cultivar, 93-11. *Theor. Appl. Genet.* 2009, *118*, 1027–1034.
- 133. Suh, J.P.; Roh, J.H.; Cho, Y.C.; Han, S.S.; Kim, Y.G.; Jena, K.K. The *Pi40* gene for durable resistance to rice blast and molecular analysis of *Pi40*-advanced backcross breeding lines. *Phytopathology* 2009, *99*, 243–250.
- Jia, Y.; Moldenhauer, K. Development of monogenic and digenic rice lines for blast resistance genes *Pi-ta*, *Pi-k^h/Pi-k^s*. J. Plant Registrations 2010, 4, 163–166.
- 135. Koide, Y.; Kawasaki, A.; Telebanco-Yanoria, M.J.; Hairmansis, A.; Nguyet, N.T.M.; Bigirimana, J.; Fujita, D.; Kobayashi, N.; Fukuta, Y. Development of pyramided lines with two resistance genes, *Pish* and *Pib*, for blast disease (*Magnaporthe oryzae*) in rice (*Oryza sativa* L.). *Plant Breed.* 2010, 129, 670–675.
- 136. Gouda, P.K.; Saikumar, S.; Varma, C.M.K.; Nagesh, K.; Thippeswamy, S.; Shenoy, V.; Ramesha, M.S.; Shashidhar, H.E. Marker-assisted breeding of *Pi-1* and *Piz-5* genes imparting resistance to rice blast in PRR78, restorer line of Pusa RH-10 Basmati rice hybrid. *Plant Breed.* 2013, *132*, 61–69.

- 137. Moumeni, A.; Leung, H. Genetic and molecular dissection of blast resistance in rice using RFLP, simple repeats and defense-related candidate gene markers. *Iran. J. Biotechnol.* **2003**, *1*, 47–54.
- 138. Abedi, F.; Babaeiyan, N.; Moumeni, A. Performance of different rice genotypes against blast pathogen through linked molecular markers. *J. Crop Sci. Biotech.* **2012**, *15*, 79–84.
- 139. Singh, V.K.; Atul, S.; Singh, S.P.; Ranjith, K.E.; Vikas, C.; Sarkel, S.; Devinder, S.; Krishnan, S.G.; Nagarajan, M.; Vinod, K.K.; *et al.* Incorporation of blast resistance into "PRR78", an elite Basmati rice restorer line, through marker assisted backcross breeding. *Field Crops Res.* 2012, *128*, 8–16.
- 140. Ashkani, S.; Rafii, M.Y.; Harun, A.R.; Latif, M.A. Genetic dissection of rice blast resistance by QTL mapping approach using an F₃ population. *Mol. Biol. Rep.* **2013**, *40*, 2503–2515.
- 141. Sallaud, C.; Lorieux, M.; Roumen, E.; Tharreau, D.; Berruyer, R.; Garsmeur, S.O.; Ghesquiere, A.; Notteghem, J.L. Identification of five new blast resistance genes in the highly blast-resistant rice variety IR64 using a QTL mapping strategy. *Theor. Appl. Genet.* **2003**, *106*, 794–803.
- 142. Kinoshita, T. Linkage mapping using mutant genes in rice. *Rice Genet Newslett.* 1998, 15, 13–74.
- 143. McCouch, S.R.; Nelson, R.J.; Thome, J.; Zeigler, R.S. *Mapping of blast resistance genes in rice*. CAB International and International Rice Research Institute; Wallingford, UK, 1994; pp. 167–187.
- 144. Yu, Z.H.; Mackill, D.J.; Bonmann, J.M.; Tanksley, S.D. Tagging genes for blast resistance in rice via linkage to RFLP markers. *Theor. Appl. Genet.* **1991**, *81*, 471–476.
- 145. Ashkani, S.; Rafii, M.Y.; Sariah, M.; Siti, N.A.A.; Rusli, I.; Harun, A.R.; Latif, M.A. Analysis of simple sequence repeat markers linked with blast disease resistance genes in a segregating population of rice (*Oryza sativa*). *Genet. Mol. Res.* 2011, 10, 1345–1355.
- 146. Correa-Victoria, F.J.; Tharreau, D.; Martinez, C.; Vales, M.; Escobar, F.; Prado, G.; Aricada, G. Combinación de genes en arroz para el desarrollo de resistencia durable a *Pyricularia grisea* en Colombia. *Fitopatol. Colomb.* 2002, *26*, 47–54.
- 147. Yamamoto, T.; Yonemaru, J.; Yano, M. Towards the understanding of complex traits in rice: Substantially or superficially? *DNA Res.* **2009**, *16*, 141–154.
- 148. Wang, Z.; Jia, Y.; Rutger, J.N.; Xia, Y. Rapid survey for presence of a blast resistance gene *Pi-ta* in rice cultivars using the dominant DNA markers derived from portions of the *Pita* gene. *Plant Breed.* **2007**, *126*, 36–42.
- 149. Liu, S.P.; Li, X.; Wang, C.Y.; Li, X.H.; He, Y.Q. Improvement of resistance to rice blast in Zhenshan 97 by molecular marker-aided selection. *Acta Bot. Sin.* **2003**, *45*, 1346–1350.
- Toojinda, T.; Tragoonrung, S.; Vanavichit, A.; Siangliw, J.L.; Pa-In, N.; Jantaboon, J.; Siangliw, M.; Fukai, S. Molecular breeding for rainfed lowland rice in the Mekong region. *Plant Prod. Sci.* 2005, *8*, 330–333.
- 151. Arnao, E.; Borges, O.; Ramis, C.; Diaz, A.; Galindo, I. Recurrent parent genome recovery in a marker assisted backcross breeding program in rice. *Interciencia* **2006**, *31*, 431–436.
- 152. Fu, C.; Wu, T.; Liu, W.; Wang, F.; Li, J.; Zhu, X.; Huang, H.; Liu, Z.R.; Liao, Y.; Zhu, M.; *et al.* Genetic improvement of resistance to blast and bacterial blight of the elite maintainer line Rongfeng B in hybrid rice (*Oryza sativa* L.) by using marker-assisted selection. *Afr. J. Biotechnol.* 2012, *11*, 13104–13114.

- 153. Miah, G.; Rafii, M.Y.; Ismail M.R.; Puteh, A.B.; Rahim, H.A.; Asfaliza, R.; Latif, M.A. Blast resistance in rice: A review of conventional breeding to molecular approaches. *Mol. Biol. Rep.* **2013**, *40*, 2369–2388.
- 154. Collard, B.C.Y.; Cruz, C.M.; McNally, K.L.; Virk, P.S.; Mackill, D.J. Rice molecular breeding laboratories in the genomics era: Current status and future and considerations. *Int. J. Plant Genomics* 2008, doi:10.1155/2008/524847.
- 155. Yu, Y.G.; Saghai-Maroof, M.A.; Buss, G.R. Divergence and allelomorphic relationship of a soybean virus resistance gene based on tightly linked DNA microsatellite and RFLP markers. *Theor. Appl. Genet.* **1996**, *92*, 64–69.
- 156. Jena, K.K.; Mackill, D.J. Molecular markers and their use in marker-assisted selection in rice. *Crop Sci.* **2008**, *48*, 1266–1276.

© 2013 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (http://creativecommons.org/licenses/by/3.0/).